


# Medical Equipment I - 2010

## Chapter 4

**Professor Yasser M. Kadah**

**Web: <http://ymk.k-space.org/courses.htm>**



# Transport in an Infinite Medium

- Definitions
- Continuity equations
- Brownian motion
- Motion in a gas
- Motion in a liquid
- Diffusion
- Applications

# Definitions

- Flow rate, volume flux or volume current ( $i$ )
  - Total volume of material transported per unit time
  - Units:  $\text{m}^3\text{s}^{-1}$
- Mass flux
- Particle flux

# Definitions

## ■ Particle fluence

- Number of particles transported per unit area across an imaginary surface
- Units:  $m^{-2}$

## ■ Volume fluence

- Volume transported per unit area across an imaginary surface
- Units:  $m^3m^{-2} = m$

# Definitions

- Fluence rate or flux density
  - Amount of “something” transported across an imaginary surface per unit area per unit time
  - Vector pointing in the direction the “something” moves and is denoted by  $\mathbf{j}$
  - Units: “something”  $\text{m}^{-2}\text{s}^{-1}$
  - Subscript to denote what “something” is

# Definitions

TABLE 4.1. Units and names for  $j$  and  $jS$  in various fields.


	$j$		$jS$	
	Units	Names	Units	Names
Particles	$\text{m}^{-2} \text{s}^{-1}$	Particle fluence rate Particle current density Particle flux density Particle flux	$\text{s}^{-1}$	Particle flux Particle current Particle flux
Electric charge	$\text{C m}^{-2} \text{s}^{-1}$ or $\text{A m}^{-2}$	Current density	$\text{C s}^{-1}$ or $\text{A}$	Current
Mass	$\text{kg m}^{-2} \text{s}^{-1}$	Mass fluence rate Mass flux density Mass flux	$\text{kg s}^{-1}$	Mass flux Mass flow
Energy	$\text{J m}^{-2} \text{s}^{-1}$ or $\text{W m}^{-2}$	Energy fluence rate Intensity Energy flux	$\text{J s}^{-1}$ or $\text{W}$	Energy flux Power

# [ Continuity Equation: 1D ]

- We deal with substances that do not “appear” or “disappear”
  - Conserved
- Conservation of mass leads to the derivation of the continuity equation

# [ Continuity Equation: 1D ]

- Consider the case of a number of particles
  - Fluence rate:  $j$  particles/unit area/unit time
- Value of  $j$  may depend on position in tube and time
  - $j = j(x, t)$
- Let volume of particles in the volume shown to be  $N(x, t)$ 
  - Change after  $\Delta t = \Delta N$





# Continuity Equation: 1D


$$\Delta N = [j(x, t) - j(x + \Delta x, t)] S \Delta t$$

- As  $\Delta x \rightarrow 0$ ,

$$j(x, t) - j(x + \Delta x, t) = -\frac{\partial j(x, t)}{\partial x} \Delta x$$


- Similarly, increase in  $N(x, t)$  is,

$$\Delta N(x, t) = N(x, t + \Delta t) - N(x, t) = \frac{\partial N}{\partial t} \Delta t$$



# Continuity Equation: 1D

- Hence,



$$\frac{\partial}{\partial t} N(x, t) = -(S\Delta x) \frac{\partial}{\partial x} j(x, t)$$

- Then, the continuity equation in 1D is,

$$\frac{\partial C}{\partial t} = - \frac{\partial j}{\partial x}$$

# Solvent Drag (Drift)

- One simple way that solute particles can move is to drift with constant velocity.
  - Carried along by the solvent,
- Process called *drift or solvent drag*.

$$\mathbf{j}_s = C \mathbf{j}_v$$

# Brownian Motion

- Application of thermal equilibrium at temperature  $T$
- Kinetic energy in 1D =  $k_B T / 2$
- Kinetic energy in 3D =  $3k_B T / 2$
- Random motion  $\rightarrow$  mean velocity  $\bar{v} = 0$ 
  - can only deal with mean-square velocity  $\overline{v^2}$

$$\frac{1}{2} m \overline{v^2} = \frac{3k_B T}{2} \Rightarrow v_{rms} = \sqrt{\overline{v^2}} = \sqrt{\frac{3k_B T}{m}}$$

# Brownian Motion

TABLE 4.2. Values of the rms velocity for various particles at body temperature.

Particle	Molecular weight	Mass (kg)	$v_{\text{rms}}$ (m s <sup>-1</sup> )
H <sub>2</sub>	2	$3.4 \times 10^{-27}$	1940
H <sub>2</sub> O	18	$3 \times 10^{-26}$	652
O <sub>2</sub>	32	$5.4 \times 10^{-26}$	487
Glucose	180	$3 \times 10^{-25}$	200
Hemoglobin	65 000	$1 \times 10^{-22}$	11
Bacteriophage	$6.2 \times 10^6$	$1 \times 10^{-20}$	1.1
Tobacco mosaic virus	$40 \times 10^6$	$6.7 \times 10^{-20}$	0.4
<i>E. coli</i>		$2 \times 10^{-15}$	0.0025

# Motion in a Gas

- Brownian motion of particles: collisions
- Mean Free Path
  - Average distance between successive collisions
- Collision Time
  - Average time between successive collisions

# Motion in a Gas

- Consider  $N(x)$  to be number of particles without collision after a distance  $x$
- For short distances  $dx$ , probability of collision is proportional to  $dx$

$$dN = N(x) \left( \frac{1}{\lambda} \right) dx$$

→

$$N(x) = N_0 e^{-x/\lambda}$$

# Motion in a Gas

- Average distance = mean free path

$$\bar{x} = \frac{1}{N_0} \int_0^{\infty} x \frac{N(x)}{\lambda} dx = -\lambda \left[ e^{-x/\lambda} \left( \frac{x}{\lambda} + 1 \right) \right]_0^{\infty} = \lambda$$

- Similar argument can be made for time


$$N(t) = N_0 e^{-t/t_c}$$

- Collision time =  $t_c$



# Motion in a Gas

- Need to evaluate  $\lambda$  and  $t_c$
- Consider one particle moving with a radius  $a_1$
- Consider stationary particles with radius  $a_2$



- Calculate collision possibilities

# Motion in a Gas

- After moving a distance  $x$ , volume covered is given by,

$$V(x) = \pi(a_1 + a_2)^2 x$$

- On average, when a particle travels mean free path, there is one collision
  - Average number of particles in  $V(\lambda)=1$
  - Concentration =  $1/V(\lambda)$

$$C = 1/V(\lambda) = 1/\pi(a_1 + a_2)^2 \lambda \Rightarrow \lambda = \frac{1}{\pi(a_1 + a_2)^2 C}$$

# Motion in a Gas

- Collision *Cross Section* is  $\pi(a_1 + a_2)^2$ 
  - Important for radiation interaction
- Example: gas at STP, volume of 1 mol = 22.4 L ( $C = 2.7 \times 10^{25} \text{m}^{-3}$ ),  $a_1 = a_2 = 0.15 \text{ nm}$ 
  - $\lambda = 0.13 \text{ } \mu\text{m}$
  - 1000 times the molecular size
  - Assumption of infrequent collisions justified

# Motion in a Gas

- Given mean free path  $\lambda$ ,

$$t_c = \frac{\lambda}{\bar{v}}$$

- Taking the average speed as  $v_{\text{rms}}$ ,

$$t_c \approx \lambda \left( \frac{m}{3k_B T} \right)^{1/2}$$


- Dependence on  $m^{1/2}$  and  $\lambda$
- For air and room temperature,  $t_c = 2 \times 10^{-10} \text{s}$

# Motion in a Liquid

- Direct substitution in Gas equations?
- For water,
  - $\lambda=a=0.1$  nm  $\rightarrow$  assumption broken
  - $t_c \sim 10^{-13}$  s  $\rightarrow$  much more frequent
  - Wrong calculations
  - However, concept appears to be valid!

# Diffusion: Fick's First Law


- Diffusion: random movement of particles from a region of higher concentration to a region of lower concentration
- Diffusing particles move independently
- Solvent at rest
  - Solute transport



# Diffusion: Fick's First Law

- If solute concentration is uniform, no net flow
- If solute concentration is different, net flow occurs

$$j_x = -D \frac{\partial C}{\partial x}$$



D: Diffusion constant ( $\text{m}^2\text{s}^{-1}$ )

# Diffusion: Fick's First Law

TABLE 4.3. Various forms of the transport equation.

Substance	Equation	Units of $j$	Units of the constant
flowing			
Particles	$j_s = -D \frac{\partial C}{\partial x}$	$\text{m}^{-2} \text{s}^{-1}$	$\text{m}^2 \text{s}^{-1}$
Mass	$j_m = -D \frac{\partial C}{\partial x}$	$\text{kg m}^{-2} \text{s}^{-1}$	$\text{m}^2 \text{s}^{-1}$
Heat	$j_H = -\kappa \frac{\partial T}{\partial x}$	$\text{J m}^{-2} \text{s}^{-1}$ or $\text{kg s}^{-3}$	$\text{J K}^{-1} \text{m}^{-1} \text{s}^{-1}$
Electric charge	$j_e = -\sigma \frac{\partial V}{\partial x}$	$\text{C m}^{-2} \text{s}^{-1}$	$\text{C m}^{-1} \text{s}^{-1} \text{V}^{-1}$ or $\Omega^{-1} \text{m}^{-1}$
Viscosity ( $y$ component of momentum transported in the $x$ direction)	$\frac{F}{S} = -\eta \frac{\partial v_y}{\partial x}$	$\text{N m}^{-2}$ or $\text{kg m}^{-1} \text{s}^{-2}$	$\text{kg m}^{-1} \text{s}^{-1}$ or $\text{Pa s}$



# Diffusion: Fick's Second Law

- Consider 1D case
- Fick's first law

$$j_x = -D \frac{\partial C}{\partial x}$$

- Continuity equation

$$\frac{\partial C}{\partial t} = -\frac{\partial j}{\partial x}$$

# Diffusion: Fick's Second Law

- Combining Fick's first law and continuity equation,

$$\frac{\partial C}{\partial t} = D \frac{\partial^2 C}{\partial x^2} \rightarrow \text{Fick's second law}$$

- 3D Case,

$$\frac{\partial C}{\partial t} = D \left( \frac{\partial^2 C}{\partial x^2} + \frac{\partial^2 C}{\partial y^2} + \frac{\partial^2 C}{\partial z^2} \right)$$


# Diffusion: Fick's Second Law

- Solving Fick's second law for  $C(x,t)$ 
  - Substitution

$$C(x,t) = \frac{N}{\sqrt{2\pi}\sigma(t)} e^{-x^2/2\sigma^2(t)}$$

where,

$$\sigma^2(t) = 2Dt + \sigma^2(0)$$



# [ Applications ]

---

- Kidney dialysis
- Tissue perfusion
- Blood oxygenation in the lung

# [ Problem Assignments ]

---

- Information posted on web site